

10. Ix JH, Allison MA, Denenberg JO, Cushman M, Criqui MH. Novel cardiovascular risk factors do not completely explain the higher prevalence of peripheral arterial disease among African Americans: The San Diego Population Study. *J Am Coll Cardiol* 2008;51:2347-54.
11. Lehto S, Niskanen L, Suhonen M, Ronnema T, Laakso M. Medial artery calcification: a neglected harbinger of cardiovascular complications in noninsulin-dependent diabetes mellitus. *Arterioscler Thromb Vasc Biol* 1996;16:978-83.
12. O'Hare AM, Katz R, Shlipak MG, Cushman M, Newman AB. Mortality and cardiovascular risk across the ankle-arm index spectrum: results from the Cardiovascular Health Study. *Circulation* 2006;113:388-93.
13. Allison MA, Hiatt WR, Hirsch AT, Coll JR, Criqui MH. A high ankle-brachial index is associated with increased cardiovascular disease morbidity and lower quality of life. *J Am Coll Cardiol* 2008;51:1292-8.
14. Allison MA, Laughlin GA, Barrett-Connor E. Association between the ankle-brachial index and carotid intimal medial thickness in the Rancho Bernardo Study. *Am J Cardiol* 2006;98:1105.
15. Allison MA, Laughlin GA, Barrett-Connor E, Langer R. Association between the ankle-brachial index and future coronary calcium (The Rancho Bernardo Study). *Am J Cardiol* 2006;97:181-6.
16. Resnick HE, Lindsay RS, McDermott MM, Devereux RB, Jones KL, Fabsitz RR, et al. Relationship of high and low ankle brachial index to all-cause and cardiovascular disease mortality: the Strong Heart Study. *Circulation* 2004;109:733-9.
17. Bild DE, Bluemke DA, Burke GL, et al. Multi-Ethnic Study of Atherosclerosis: objectives and design. *Am J Epidemiol* 2002;156:871-81.
18. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972;18:499-502.
19. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2004; 27:S5-10.
20. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* 2002;39:S1-266.
21. McDermott MM, Greenland P, Liu K, Guralnik JM, Celis L, Criqui MH, et al. The ankle-brachial index is associated with leg function and physical activity: the Walking and Leg Circulation Study. *Ann Intern Med* 2002;136:873-83.
22. Glymour MM, Weuve J, Berkman LF, Kawachi I, Robins JM. When is baseline adjustment useful in analyses of change? An example with education and cognitive change. *Am J Epidemiol* 2005;162:267-78.
23. Yanez ND III, Kronmal RA, Shemanski LR, Psaty BM. A regression model for longitudinal change in the presence of measurement error. *Ann Epidemiol* 2002;12:34-8.
24. Ouriel K, Zarins CK. Doppler ankle pressure: an evaluation of three methods of expression. *Arch Surg* 1982;117:1297-300.
25. Fowkes FG. Epidemiology of peripheral vascular disease. *Atherosclerosis* 1997;131 Suppl:S29-31.
26. Spittell JA Jr. Peripheral arterial disease. *Dis Mon* 1994;40:641-700.
27. Brooks B, Dean R, Patel S, Wu B, Molyneaux L, Yue DK. TBI or not TBI: that is the question. Is it better to measure toe pressure than ankle pressure in diabetic patients? *Diabet Med* 2001;18:528-32.
28. Aboyans V, Criqui MH, McClelland RL, Allison MA, McDermott MM, Goff DC, et al. Intrinsic contribution of gender and ethnicity to normal ankle-brachial index values: the Multi-Ethnic Study of Atherosclerosis (MESA). *J Vasc Surg* 2007;45:319-27.
29. Shadman R, Criqui MH, Bundens WP, Fronck A, Denenberg JO, Gamst AC, et al. Subclavian artery stenosis: prevalence, risk factors, and association with cardiovascular diseases. *J Am Coll Cardiol* 2004;44: 618-23.
30. Din-Dzietham R, Couper D, Evans G, Arnett DK, Jones DW. Arterial stiffness is greater in African Americans than in whites: evidence from the Forsyth County, North Carolina, ARIC cohort. *Am J Hypertens* 2004;17:304-13.
31. Libby P. Vascular biology of atherosclerosis: overview and state of the art. *Am J Cardiol* 2003;91:3A-6A.
32. Aboyans V, Ho E, Denenberg JO, Ho LA, Natarajan L, Criqui MH. The association between elevated ankle systolic pressures and peripheral occlusive arterial disease in diabetic and nondiabetic subjects. *J Vasc Surg* 2008;48:1197-203.
33. Aboyans V, Criqui MH, Denenberg JO, Knoke JD, Ridker PM, Fronck A. Risk factors for progression of peripheral arterial disease in large and small vessels. *Circulation* 2006;113:2623-9.
34. Meijer WT, Grobbee DE, Hunink MGM, Hofman A, Hoes AW. Determinants of peripheral arterial disease in the elderly: the Rotterdam Study. *Arch Intern Med* 2000;160:2934-8.

Submitted Mar 3, 2009; accepted May 6, 2009.

## INVITED COMMENTARY

Carlos H. Timaran, MD, Dallas, Tex

Changes in the ankle-brachial index (ABI) and progression to peripheral arterial disease (PAD) may result in major detrimental effects on quality of life and functional status and, more importantly, are potent predictors of myocardial infarction, stroke, and death. Despite its frequent occurrence, little is known about the natural history of PAD among minority ethnic groups.

The Multi-Ethnic Study of Atherosclerosis (MESA) was designed to determine ethnic differences in the progression of subclinical to clinical cardiovascular disease.

In the current study, Allison et al assessed changes in ABI in the MESA cohort over a 3-year time period. The authors conclude that besides traditional risk factors, African Americans had a significantly higher risk of progression to a low ABI ( $<0.9$ ) and a reduced trend towards progression to a high ABI ( $>1.4$ ). Conversely, Hispanics and Chinese did not reveal significant changes in ABI over time. These findings should not come as a surprise as African Americans have repeatedly been shown to have a higher prevalence of PAD, which appears independent of both traditional and novel vascular risk factors. There are, however, several uncertainties and ambiguities that deserve further investigation.

Although PAD has traditionally been defined as resting ABI of less than 0.90, the ABI may underestimate the severity of disease in individuals with calcified, noncompressible arteries, which are more frequent in individuals with diabetes, chronic renal failure, or

advanced age. Although the literature quoted by most epidemiologic studies, including this one, regarding the accuracy of the ABI to detect PAD is primarily based in small clinical observational studies that assessed individuals with established PAD prior to treatment, the ABI has been accepted without adequate scrutiny in all populations, including those with high prevalence of diabetes. In the MESA cohort, Hispanics did not have an increased risk of significant ABI changes despite similar age and sex-adjusted prevalence of baseline diabetes and dyslipidemia compared with African Americans. Allison et al claim that the stability of ABI among Hispanics may be related to either no actual changes or similar progression in atherosclerosis and arterial rigidity. If one assumes that, as suggested by the authors, progression is the rule, the lack of changes in ABI may be more likely related to progression of both atherosclerosis and arterial rigidity. It may also mean that among Hispanics, there may be individuals with either baseline or subsequent stenosis of 50% or more in a major leg artery, ie, PAD, that may not result in lower ABI because of the "compensatory" increased arterial rigidity. The ABI may, therefore, be inaccurate to detect PAD in Hispanics given their high prevalence of diabetes and vascular calcification that render the ABI unreliable. Other noninvasive tests, such as pulse volume recordings and toe-brachial index, in addition to the ABI, should be assessed for accurate detection and screening of PAD in this defined population.